

Equilibria of Deferred Acceptance with Complete Lists*

Bettina Klaus[†] Flip Klijn[‡]

April 1, 2016

Abstract

We study the structure of the set of (Nash) equilibria of a deferred acceptance game with complete lists: for a given marriage market with complete lists, men propose to women truthfully while women can accept or reject proposals strategically throughout the deferred-acceptance algorithm. Zhou (1991) studied this game and showed that a matching that is stable with respect to the true preferences can be supported by some preference profile (possibly a non-equilibrium one) if and only if it can be supported by an equilibrium as well. In particular, this result implies the existence of equilibria since the men-optimal stable matching is supported by true preferences and hence an equilibrium outcome. We answer an open question Zhou posed by showing that there need not exist an equilibrium matching that weakly dominates all other equilibrium matchings from the women’s point of view (Theorem 2). We complement Zhou’s and our findings by showing that the set of equilibrium matchings also need not be “connected” (Example 2).

Keywords: matching, stability, complete lists, Nash equilibria.

JEL-Numbers: C72, C78, D47.

*We thank Battal Doğan for comments on a previous version.

[†]Faculty of Business and Economics, University of Lausanne, Internef 538, CH-1015 Lausanne, Switzerland; e-mail: bettina.klaus@unil.ch. B. Klaus gratefully acknowledges financial support from the Swiss National Science Foundation (SNFS).

[‡]*Corresponding author.* Institute for Economic Analysis (CSIC) and Barcelona GSE, Campus UAB, 08193 Bellaterra (Barcelona), Spain; e-mail: flip.klijn@iae.csic.es. F. Klijn gratefully acknowledges financial support from the Generalitat de Catalunya (2014-SGR-1064), the Spanish Ministry of Economy and Competitiveness through Plan Nacional I+D+i (ECO2014-59302-P), and the Severo Ochoa Programme for Centres of Excellence in R&D (SEV-2015-0563).

1 Introduction

The so-called marriage model is concerned with (two-sided, one-to-one) matching markets where the two sides of the market are, for instance, men and women (or firms and workers). A matching is then a partition of all men and women into couples and unmatched agents. Such a matching is “stable” if each man and woman has an acceptable match, and no man and woman prefer one another to their respective matches. Gale and Shapley (1962) were the first to formalize this notion of stability and presented their deferred-acceptance algorithm to calculate stable matchings. Stability proved to be an essential property in several real-life markets. For instance, in many centralized labor markets, clearinghouses are most often successful if they produce stable matchings.¹ This inspired many researchers to study stability; see Roth and Sotomayor (1990) for a classic survey, and Vulkan et al. (2013) and Manlove (2013) for more recent surveys.

In the original Gale-Shapley model, the preference lists of the agents are assumed to be complete, *i.e.*, for each agent all agents on the other side are acceptable. In other words, remaining unmatched is not desirable. Moreover, agents cannot be (falsely) declared unacceptable. The men-proposing deferred-acceptance algorithm with complete lists induces a game where it is a weakly dominant strategy for each man to reveal his preferences truthfully (Dubins and Freedman, 1981; Roth, 1982). For marriage markets with complete lists, Roth (1984b) showed that the matching induced by any equilibrium in (weakly) undominated strategies is stable with respect to the true preferences. Zhou (1991) assumed that men propose to women truthfully while women can accept or reject proposals strategically throughout the deferred-acceptance algorithm and showed that a matching that is stable with respect to the true preferences can be supported by some preference profile (possibly a non-equilibrium one) if and only if it can be supported by an equilibrium as well. In particular, this result implies the existence of equilibria since the men-optimal stable matching is supported by true preferences and hence an equilibrium outcome. Zhou (1991) conjectured the existence of an equilibrium matching (possibly not the women-optimal stable matching) that weakly dominates all other equilibrium matchings from the women’s point of view. Teo et al. (2001) showed that it is not always possible for a woman w to obtain her women-optimal stable partner from the men-proposing deferred-acceptance algorithm, but they assumed that woman w is the only agent acting strategically, *i.e.*, their study does not consider the equilibria of the deferred-acceptance game.

In this note, we solve Zhou’s (1991) conjecture in the negative by showing that there need not exist an equilibrium matching (the women-optimal stable matching or not) that weakly dominates all other equilibrium matchings from the women’s point of view (Theorem 2). We complement Zhou’s and our findings by showing that the set of equilibrium matchings also need not be “connected.” More precisely, suppose two stable matchings are supported by equilibria. Then, it can happen that any stable matching that is located between the two stable matchings (in terms of each agent’s preferences) is not supported

¹See Roth (1984a) and Roth and Xing (1994) for empirical evidence.

by an equilibrium (Example 2).

The results for the original Gale-Shapley model contrast with those for the model that allows for “rejections,” *i.e.*, an agent’s preference list need not be complete in the sense that some agents on the other side of the market may be unacceptable. In this case, agents have the option of declaring some agents as unacceptable. In the game induced by the men-proposing deferred-acceptance algorithm, the set of equilibrium outcomes coincides with the set of stable matchings (Gale and Sotomayor, 1985; Roth, 1984b). An important consequence is that the remarkable properties of the set of stable matchings carry over to the set of equilibrium outcomes. In particular, there exist an equilibrium matching (the women-optimal stable matching) that weakly dominates all other equilibrium matchings from the women’s point of view. However, for the more general many-to-one matching model such an equilibrium matching also does not need to exist (Jaramillo et al., 2013, Example 1).

2 Model

There are two finite and disjoint sets of agents: a set $M = \{m_1, \dots, m_n\}$ of men and a set $W = \{w_1, \dots, w_n\}$ of women. Thus, $|M \cup W| = 2n$. We denote a generic agent by i , a generic man by m , and a generic woman by w .

Each agent has a complete, transitive, and strict preference relation over the agents on the other side of the market. Hence, man m ’s preferences \succ_m can be represented as a strict ordering P_m of the elements in W , for instance: $P_m = w_3 w_4 w_2 w_1$ which indicates that m prefers w_3 to w_4 to w_2 to w_1 . Similarly, woman w ’s preferences \succ_w can be represented as a strict ordering P_w of the elements in M . For any $i \in M \cup W$, we write $j \succeq_i k$ if $j \succ_i k$ or $j = k$. For any $I \subseteq M \cup W$, we define $P_I \equiv (P_i)_{i \in I}$. We write P instead of $P_{M \cup W}$.

A **matching market** is a triple (M, W, P) , or shortly P . A matching for (M, W, P) is a function $\mu : M \cup W \rightarrow M \cup W$ such that for all $m \in M$ and $w \in W$ it holds that $\mu(m) = w \Leftrightarrow \mu(w) = m$. If $\mu(m) = w$, then man m and woman w are matched to one another. We call $\mu(i)$ the match of agent i at μ . When denoting a matching μ we list the women that are matched to men m_1, \dots, m_n ; e.g., $\mu = w_3, w_4, w_2, w_1$ denotes a matching where m_1 is matched to w_3 , m_2 to w_4 , m_3 to w_2 , and m_4 to w_1 . Alternatively, a matching can be described by the list of men that are matched to women w_1, \dots, w_n .

A key property of matchings is stability. If an agent can improve upon its present match by switching to another agent such that this agent is better off as well, then this blocking clearly would cause instability. For a given matching μ , a man m and a woman w are a blocking pair if they are not matched to one another but prefer one another to their current match at μ , *i.e.*, $w \succ_m \mu(m)$ and $m \succ_w \mu(w)$. A matching is **stable** if there are no blocking pairs. We denote the set of stable matchings for matching market P by $\mathcal{S}(P)$. Gale and Shapley (1962) proved that $\mathcal{S}(P) \neq \emptyset$.

In fact, Gale and Shapley (1962) showed that there is a **men-optimal (women-pessimal) stable matching** μ_M^P , *i.e.*, for each man $m \in M$, each woman $w \in W$,

and each stable matching μ , $\mu_M^P(m) \succeq_m \mu(m)$ and $\mu(w) \succeq_w \mu_M^P(w)$. Similarly, there is a **men-pessimal (women-optimal) stable matching** μ_W^P . Gale and Shapley (1962) provided an algorithm, called the deferred-acceptance algorithm, to compute μ_M^P and μ_W^P . Next, we describe the (men-proposing) deferred-acceptance (DA) algorithm² to obtain the men-optimal stable matching $\mu_M^P \in \mathcal{S}(P)$ for any preference profile P .

DA algorithm:

STEP 1. Each man proposes to his most preferred woman. Each woman who receives at least one proposal is tentatively matched to her most preferred proposer and rejects all other proposers.

STEP $k \geq 2$. Each man who has been rejected in Step $k - 1$ proposes to his most preferred woman among the ones that have not rejected him yet. Each woman who receives at least one proposal is tentatively matched to her most preferred man among the ones that proposed to her and the one she is currently tentatively matched to (if any)– all other proposers are rejected.

The algorithm terminates when no man is rejected. Then, tentative matches become final, and the resulting matching is μ_M^P . Let φ be the function that associates each preference profile P with the men-optimal stable matching μ_M^P for P , *i.e.*, $\varphi(P) = \mu_M^P$.

The DA algorithm induces a game form where each agent can reveal a preference relation over the other side of the market. For each strategy profile Q , the DA algorithm produces $\varphi(Q)$ as the outcome. With each true preference profile P , the DA algorithm induces a **game** $\Gamma(P)$. Dubins and Freedman (1981) and Roth (1982) showed that for each man $m \in M$ it is a weakly dominant strategy to play P_m , *i.e.*, to reveal his preferences truthfully. Henceforth, we will assume that *men always state the truth* and that *women are the only strategic agents*.

Roth (1984b) showed that for any strategy profile Q that is a Nash equilibrium (in weakly undominated strategies) of the game $\Gamma(P)$, matching $\varphi(Q)$ is stable with respect to the true preferences P . Zhou (1991) proved the existence of Nash equilibria of $\Gamma(P)$ and characterized the matchings that can be sustained at Nash equilibria. Let $\mathcal{E}(P)$ be the matchings that are the outcome of some Nash equilibrium of $\Gamma(P)$. Let $\mathcal{O}(P_M)$ be the matchings that can be obtained at some strategy profile where men state the truth and women play any set of strategies (which do not necessarily constitute a Nash equilibrium), *i.e.*,

$$\mathcal{O}(P_M) = \{\varphi(P_M, Q_W) : Q_W \text{ is a strategy profile of the women}\}.$$

Theorem 1. [Zhou, 1991, Theorem 1]

For each market P , $\mathcal{E}(P) = \mathcal{S}(P) \cap \mathcal{O}(P_M)$. In particular, since $\mu_M^P = \varphi(P_M, P_W) \in \mathcal{S}(P)$, $\mathcal{E}(P) \neq \emptyset$.

Zhou (1991, Theorem 2) also showed that $\mathcal{E}(P)$ contains matchings different from μ_M^P if truth-telling is not an equilibrium profile for women.

²By switching the roles of men and women in the deferred-acceptance algorithm, matching μ_W^P is obtained.

3 Results

Zhou (1991, p. 29) asked whether women can coordinate their manipulations in an optimal fashion. More precisely, “does there exist a matching [in] $\mathcal{E}(P)$ that weakly dominates all other matchings in $\mathcal{E}(P)$ from the women’s point of view? Such a matching, if [it] exists, seems more likely to emerge than others. . . . We leave it as a conjecture for future research.”

The next theorem answers Zhou’s (1991) question.

Theorem 2. *If $n \leq 3$, then for any market P there exists a matching $\mu \in \mathcal{E}(P)$ that weakly dominates all other matchings $\nu \in \mathcal{E}(P)$ from the women’s point of view, i.e., for all $w \in W$, $\mu(w) \succeq_w \nu(w)$. If $n \geq 4$, this is not necessarily true.*

Proof. Recall that by Theorem 1, for all markets P , $\mathcal{E}(P) \subseteq \mathcal{S}(P)$.

Let $n \leq 2$. Then, for any market P there are at most 2 stable matchings, and hence the statement is trivially true.

Let $n = 3$. Let P be a market. Suppose by contradiction that there exists no matching in $\mathcal{E}(P)$ that weakly dominates all other matchings in $\mathcal{E}(P)$ from the women’s point of view. Then, women cannot find a jointly optimal equilibrium matching and hence there exist at least two distinct (stable) matchings $\mu', \mu'' \in \mathcal{E}(P)$, each of which is considered optimal by some women but not others. Thus, there are distinct women, say w_1 and w_2 , such that woman w_1 finds μ' optimal but not μ'' [for all $\nu \in \mathcal{E}(P)$, $\mu'(w_1) \succeq_{w_1} \nu(w_1)$ and $\mu'(w_1) \succ_{w_1} \mu''(w_1)$] while woman w_2 finds μ'' optimal but not μ' [for all $\nu \in \mathcal{E}(P)$, $\mu''(w_2) \succeq_{w_2} \nu(w_2)$ and $\mu''(w_2) \succ_{w_2} \mu'(w_2)$].

Let $W(\mu')$ be the set of women who strictly prefer μ' to μ'' and $M(\mu')$ be the set of men who strictly prefer μ' to μ'' . Analogously define $W(\mu'')$ and $M(\mu'')$. Since $\mu', \mu'' \in \mathcal{E}(P) \subseteq \mathcal{S}(P)$ it follows from Donald Knuth’s decomposition lemma (Roth and Sotomayor, 1990, Corollary 2.21) that for all $w \in W$, [$w \in W(\mu')$ if and only if $\mu'(w) \in M(\mu'')$] and [$w \in W(\mu'')$ if and only if $\mu''(w) \in M(\mu')$].

Note that $w_1 \in W(\mu')$ and $w_2 \in W(\mu'')$. Then, $w_1 \in W(\mu')$ implies $m' \equiv \mu'(w_1) \in M(\mu'')$. By assumption, w_1 gets different matches at μ' and μ'' . Hence, $m' = \mu'(w_1)$ gets different matches at μ' and μ'' . But then also $w'' \equiv \mu''(m')$ gets different matches at μ' and μ'' . Hence, $w'' \in W(\mu') \cup W(\mu'')$. Suppose $w'' \in W(\mu'')$. Since $m' \in M(\mu'')$ and $w'' = \mu''(m')$, (m', w'') is a blocking pair for μ' , which contradicts the stability of μ' . Hence, $w'' \in W(\mu')$. Since (i) $\mu'(m') = w_1$ and $\mu''(m') = w''$ and (ii) m' gets different matches at μ' and μ'' , it follows that $w_1 \neq w''$. Hence, $|W(\mu')| \geq 2$. Similarly, it follows that $w_2 \in W(\mu'')$ implies $|W(\mu'')| \geq 2$. Since $W(\mu') \cap W(\mu'') = \emptyset$, it follows that $|W| \geq |W(\mu')| + |W(\mu'')| \geq 4$, which contradicts $|W| = n = 3$.

If $n = 4$, then Example 1 shows that there need not exist an equilibrium matching that weakly dominates all other equilibrium matchings from the women’s point of view. The example extends to the case $n > 4$ by making for each $l > 4$, the members of (m_l, w_l) each other’s mutually best possible partner. \square

Example 1. [No optimal equilibrium outcome for the women]

Consider the matching market (M, W, P) where $M = \{m_1, m_2, m_3, m_4\}$, $W = \{w_1, w_2, w_3, w_4\}$, and preferences P given by Table 1. In each column, higher placed agents are more preferred agents.

men				women			
m_1	m_2	m_3	m_4	w_1	w_2	w_3	w_4
w_1	w_4	w_1	w_4	m_2	m_1	m_4	m_3
w_2	w_2	w_3	w_3	m_1	m_2	m_3	m_4
\dots	w_1	w_4	\dots	\dots	\dots	\dots	\dots
	w_3	w_3					

Table 1: Preferences P in Example 1

The entries \dots can be any agents as long as each column is a preference relation over the agents on the other side of the market. The set of stable matchings, given by $\mathcal{S}(P) = \{\mu_M^P = \mu_1, \mu_2, \mu_3, \mu_4 = \mu_W^P\}$, is depicted in Table 2.³ In all tables, the men-optimal stable matching is depicted in boldface while the women-optimal stable matching is the boxed matching.

	men				women			
	m_1	m_2	m_3	m_4	w_1	w_2	w_3	w_4
$\mu_M^P = \mu_1 :$	w_1	w_2	w_3	w_4	m_1	m_2	m_3	m_4
$\mu_2 :$	w_2	w_1	w_3	w_4	m_2	m_1	m_3	m_4
$\mu_3 :$	w_1	w_2	w_4	w_3	m_1	m_2	m_4	m_3
$\mu_W^P = \mu_4 :$	w_2	w_1	w_4	w_3	m_2	m_1	m_4	m_3

Table 2: The four stable matchings in Example 1

To show that there is no optimal equilibrium outcome for the women it is sufficient to prove that $\mathcal{E}(P) = \{\mu_1, \mu_2, \mu_3\}$: for all women $w \in \{w_1, w_2\}$, $\mu_2(w) \succ_w \mu_1(w) = \mu_3(w)$, while for all women $w \in \{w_3, w_4\}$, $\mu_3(w) \succ_w \mu_1(w) = \mu_2(w)$.

We first show that $\mathcal{E}(P) \supseteq \{\mu_1, \mu_2, \mu_3\}$. By Theorem 1, $\mu_1 = \mu_M^P \in \mathcal{E}(P)$. Next, note that for profile $Q^2 \equiv (P_M, Q_W^2)$, where Q_W^2 are the women's preferences given in Table 3, we have $\varphi(Q^2) = \mu_2$. Hence, $\mu_2 \in \mathcal{O}(P_M)$. So, by Theorem 1, $\mu_2 \in \mathcal{E}(P)$. Similarly, for profile $Q^3 \equiv (P_M, Q_W^3)$, where Q_W^3 are the women's preferences given in Table 4, we have $\varphi(Q^3) = \mu_3$, and hence, $\mu_3 \in \mathcal{E}(P)$.

³It is easy to see that the set of stable matchings is indeed $\mathcal{S}(P) = \{\mu_M^P = \mu_1, \mu_2, \mu_3, \mu_4 = \mu_W^P\}$. First, one computes μ_M^P and μ_W^P using the two versions of the DA algorithm. Second, one verifies that the only other two matchings μ with $\mu_M^P(m) \succeq_m \mu(m) \succeq_m \mu_W^P(m)$ for all $m \in M$ are μ_2 and μ_3 . Finally, one checks the stability of μ_2 and μ_3 .

women			
w_1	w_2	w_3	w_4
m_2	m_1	m_4	m_3
m_3	m_2	m_3	m_4
m_1	\dots	\dots	\dots
m_4			

Table 3: Preferences Q_W^2

women			
w_1	w_2	w_3	w_4
m_2	m_1	m_4	m_3
m_1	m_2	m_3	m_2
\dots	\dots	\dots	m_4
			m_1

Table 4: Preferences Q_W^3

Next, we show that $\mathcal{E}(P) \subseteq \{\mu_1, \mu_2, \mu_3\}$. By Theorem 1, $\mathcal{E}(P) = \mathcal{S}(P) \cap \mathcal{O}(P_M) \subseteq \{\mu_1, \mu_2, \mu_3, \mu_4\}$. Hence, it suffices to show that $\mu_4 \notin \mathcal{E}(P)$. Since $\mu_4 \in \mathcal{S}(P)$, we have to prove that $\mu_4 \notin \mathcal{O}(P_M)$, *i.e.*, for all possible preference profiles Q_W of the women, $\varphi(P_M, Q_W) \neq \mu_4$. Let Q_W be a preference profile of the women. Notice that in Step 1 of the DA algorithm, men m_1 and m_3 propose to woman w_1 and men m_2 and m_4 propose to woman w_4 . Since each of these women has to reject one applicant, we distinguish between the four cases⁴ depicted in the first column of Table 5. In all cases, the DA algorithm terminates in Step 2, *i.e.*, the other two women receive exactly one proposal that they have to accept.

Case	men				women				
	m_1	m_2	m_3	m_4	w_1	w_2	w_3	w_4	
(a) $m_1 Q_{w_1} m_3$ and $m_4 Q_{w_4} m_2$	$\Rightarrow \mu_1 :$	w_1	w_2	w_3	w_4	m_1	m_2	m_3	m_4
(b) $m_3 Q_{w_1} m_1$ and $m_4 Q_{w_4} m_2$	$\Rightarrow \mu_2 :$	w_2	w_1	w_3	w_4	m_2	m_1	m_3	m_4
(c) $m_1 Q_{w_1} m_3$ and $m_2 Q_{w_4} m_4$	$\Rightarrow \mu_3 :$	w_1	w_2	w_4	w_3	m_1	m_2	m_4	m_3
(d) $m_3 Q_{w_1} m_1$ and $m_2 Q_{w_4} m_4$	$\Rightarrow \mu_5 :$	w_2	w_4	w_1	w_3	m_3	m_1	m_4	m_2

Table 5: The four outcomes $\varphi(P_M, Q_W)$ in Example 1

By applying the men-proposing DA algorithm, one easily verifies that in each case the resulting matching $\varphi(P_M, Q_W)$ is the one denoted in the same row. Since in all four cases $\varphi(P_M, Q_W) \neq \mu_4$ (note that matching μ_5 is not stable), the proof is completed. \diamond

Our Example 1 and Example 1 in Zhou (1991) may suggest that for any market P the set of equilibria is “connected,” *i.e.*, if $\mu, \mu' \in \mathcal{E}(P)$, then for any $\nu \in \mathcal{S}(P)$ with $\mu(m) \succeq_m \nu(m) \succeq_m \mu'(m)$ for all $m \in M$, $\nu \in \mathcal{E}(P)$. The following example shows this need not be the case.

⁴For instance, $m_1 Q_{w_1} m_3$ (in Cases (a) and (c)) means that in w_1 's strategy (list) man m_1 is more preferred to m_3 and hence she rejects m_3 at Step 1 of the DA algorithm.

Example 2. [Gap in set of equilibrium outcomes]

Consider the matching market (M, W, P) where $M = \{m_1, m_2, m_3, m_4, m_5, m_6\}$, $W = \{w_1, w_2, w_3, w_4, w_5, w_6\}$, and preferences P given by Table 6.

men						women					
m_1	m_2	m_3	m_4	m_5	m_6	w_1	w_2	w_3	w_4	w_5	w_6
<u>w₂</u>	w₄	w₅	w₃	<u>w₁</u>	w_1	m_1	m_2	m_3	m_4	m_5	m_6
w_1	w_3	w_4	w_5	w_5	w₆	m_5	m_1	m_2	m_3	m_4	...
...	w_2	w_3	w_4	m_6	...	m_4	m_2	m_3	...
	

Table 6: Preferences P in Example 2

The entries ... can again be any agents as long as each column is a preference relation over the agents on the other side of the market. The set of stable matchings, given by $\mathcal{S}(P) = \{\nu_1, \nu_2, \nu_3\}$, is depicted in Table 7.⁵ In all tables, the men-optimal stable matching ν_1 is depicted in boldface, the women-optimal stable matching ν_3 is the boxed matching, and the only other stable matching ν_2 is underlined.

	men						women					
	m_1	m_2	m_3	m_4	m_5	m_6	w_1	w_2	w_3	w_4	w_5	w_6
ν_1	w₂	w₄	w₅	w₃	w₁	w₆	m_5	m_1	m_4	m_2	m_3	m_6
ν_2	<u>w₂</u>	<u>w₃</u>	<u>w₄</u>	<u>w₅</u>	<u>w₁</u>	<u>w₆</u>	<u>m_5</u>	<u>m_1</u>	<u>m_2</u>	<u>m_3</u>	<u>m_4</u>	<u>m_6</u>
ν_3	w_1	w_2	w_3	w_4	w_5	w_6	m_1	m_2	m_3	m_4	m_5	m_6

Table 7: The three stable matchings in Example 2

We will show that $\mathcal{E}(P) = \{\nu_1, \nu_3\}$. We first show that $\mathcal{E}(P) \supseteq \{\nu_1, \nu_3\}$. By Theorem 1, $\nu_1 \in \mathcal{E}(P)$. Next, note that for profile $\bar{Q}^3 \equiv (P_M, \bar{Q}_W^3)$, where \bar{Q}_W^3 are the women's preferences given in Table 8, we have $\varphi(\bar{Q}^3) = \nu_3$. Hence, $\nu_3 \in \mathcal{O}(P_M)$. So, by Theorem 1, $\nu_3 \in \mathcal{E}(P)$.

Next, we show that $\mathcal{E}(P) \subseteq \{\nu_1, \nu_3\}$. By Theorem 1, $\mathcal{E}(P) = \mathcal{S}(P) \cap \mathcal{O}(P_M) \subseteq \{\nu_1, \nu_2, \nu_3\}$. Hence, it suffices to show that $\nu_2 \notin \mathcal{E}(P)$. Suppose to the contrary that there is a preference profile Q_W of the women such that (P_M, Q_W) is an equilibrium and $\varphi(P_M, Q_W) = \nu_2$. Notice that in Step 1 of the DA algorithm, men m_5 and m_6 propose to woman w_1 . Hence, for women w_1 we can distinguish between two cases.

Suppose (a) $m_5 Q_{w_1} m_6$. Then, in woman w_1 's strategy (list) man m_5 is more preferred to m_6 and hence she rejects m_6 at Step 1 of the DA algorithm. Furthermore, the DA

⁵It is easy to see that the set of stable matchings is indeed $\mathcal{S}(P) = \{\nu_1, \nu_2, \nu_3\}$. First, the two versions of the DA algorithm yield ν_1 and ν_3 . Second, one verifies that among all matchings ν that satisfy $\nu_1(m) \succeq_m \nu(m) \succeq_m \nu_3(m)$ for all $m \in M$ the only other stable matching is ν_2 .

women					
w_1	w_2	w_3	w_4	w_5	w_6
m_1	m_2	m_3	m_4	m_5	m_6
m_6	m_1	m_4	m_2	m_3	\dots
m_5	\dots	m_2	m_3	\dots	
\dots		\dots	\dots		

Table 8: Preferences \bar{Q}_W^3

algorithm terminates in Step 2, *i.e.*, all other women receive exactly one proposal that they have to accept and $\varphi(P_M, Q_W) = \nu_1 \neq \nu_2$.

Now suppose (b) $m_6 Q_{w_1} m_5$. Then, at Step 1 of the DA algorithm, woman w_1 rejects m_5 , who at Step 2 proposes to w_5 . Since m_5 is woman w_5 's most preferred man (according to her true preferences P_{w_5}) and since (P_M, Q_W) is an equilibrium, w_5 and m_5 are matched to one another at $\varphi(P_M, Q_W)$, but then $\varphi(P_M, Q_W) \neq \nu_2$. Hence, $\nu_2 \notin \mathcal{E}(P)$. \diamond

References

- Dubins, L. E. and Freedman, D. A. (1981): “Machiavelli and the Gale-Shapley Algorithm.” *American Mathematical Monthly*, 88: 485–494.
- Gale, D. and Shapley, L. S. (1962): “College Admissions and the Stability of Marriage.” *American Mathematical Monthly*, 69: 9–15.
- Gale, D. and Sotomayor, M. A. O. (1985): “Ms Machiavelli and the Stable Matching Problem.” *American Mathematical Monthly*, 92: 261–268.
- Jaramillo, P., Kayı, Ç., and Klijn, F. (2013): “Equilibria under Deferred Acceptance: Dropping Strategies, Filled Positions, and Welfare.” *Games and Economic Behavior*, 82: 693–701.
- Manlove, D. (2013): *Algorithmics of Matching under Preferences*. World Scientific Publishing.
- Roth, A. E. (1982): “The Economics of Matching: Stability and Incentives.” *Mathematics of Operations Research*, 7: 617–628.
- Roth, A. E. (1984a): “The Evolution of the Labor Market for Medical Interns and Residents: A Case Study in Game Theory.” *Journal of Political Economy*, 92: 991–1016.
- Roth, A. E. (1984b): “Misrepresentation and Stability in the Marriage Problem.” *Journal of Economic Theory*, 34: 383–387.
- Roth, A. E. and Sotomayor, M. A. O. (1990): *Two-Sided Matching: A Study in Game-Theoretic Modeling and Analysis*. Cambridge University Press, Cambridge.

- Roth, A. E. and Xing, X. (1994): “Jumping the Gun: Imperfections and Institutions related to the Timing of Market Transactions.” *American Economic Review*, 84: 992–1044.
- Teo, C. P., Sethuraman, J., and Tan, W. P. (2001): “Gale-Shapley Stable Marriage Problem Revisited: Strategic Issues and Applications.” *Management Science*, 47: 1252–1267.
- Vulkan, N., Roth, A. E., and Neeman, Z., editors (2013): *The Handbook of Market Design*. Oxford University Press.
- Zhou, L. (1991): “Stable Matchings and Equilibrium Outcomes of the Gale–Shapley’s Algorithm for the Marriage Problem.” *Economics Letters*, 36: 25–29.