

# Abstract

The Omitiomire Cu deposit in the Ekuja Dome of the Damara Belt in Namibia is hosted by an anastomosing, low-angle Pan-African (ca. 520 Ma, Miller 1983, 2008; Gray et al. 2008.) shear zone system developed around an older (ca. 1100–1060 Ma, Steven et al. 2000; Maiden et al., 2014), late Mesoproterozoic intrusive breccia between a suite of mafic rocks (originally lava flows) and later felsic gneisses. High-grade ore copper is formed along contacts between tectonically interleaved biotite-epidote schists and felsic gneisses, and are in contact with metabasite.

Alteration and mineralization are associated with elevated concentrations of K<sub>2</sub>O, and Cu and a loss of Na<sub>2</sub>O, CaO, and MgO. Oxygen isotope fractionation for quartz-biotite, quartz-feldspar, and quartz amphibole mineral pairs give temperatures of between 500 and 650 °C during peak of metamorphism. Mineral separates from amphibole-biotite gneisses and mineralized schists have similar ranges in  $\delta^{18}\text{O}$  values of about 1.2 to 2 ‰ relative to VSMOW.

Coexisting minerals are arranged in an order of increasing  $\delta^{18}\text{O}$  values from biotite, to epidote, amphibole, and quartz, suggesting that was balance with the rock and the chemical composition of the silicates. Similarly, H-isotope results for mineral separates from biotite-epidote schists and amphibole gneisses do not show any reversals for D/H fractionations, with  $\delta\text{D}$  values of between -46 and -82 ‰, typical of metamorphic-magmatic rocks.

The homogeneous and low  $\delta^{34}\text{S}$  values (-6.2 to -4.5‰ CDT) are compatible with a local redistribution of sulfur from magmatic rocks and interaction with sulfur derived from metamorphic fluids during Kuiseb schist alteration.

The relatively low fluid/rock ratios and elevated Cu values within amphibolite such as a metabasalt or a meta andesite point to a local redistribution of Cu mineralization. The magmatic origin of a copper enriched metabasite with hydrothermal alteration seem to be related to a metamorphosed Volcanic Mafic sulfide. (Bottinga Y., 1975)